# The Six-Point Circle Theorem \*

## Adrian Mitrea

#### Abstract

Given  $\triangle ABC$  and angles  $\alpha, \beta, \gamma \in (0, \pi)$  with  $\alpha + \beta + \gamma = \pi$ , we study the properties of the triangle DEF which satisfies: (i)  $D \in BC$ ,  $E \in AC$ ,  $F \in AB$ , (ii)  $\not\Delta D = \alpha$ ,  $\not\Delta E = \beta$ ,  $\not\Delta F = \gamma$ , (iii)  $\Delta DEF$  has the minimal area in the class of triangles satisfying (i) and (ii). In particular, we show that minimizer  $\Delta DEF$ , exists, is unique and is a pedal triangle, corresponding to a certain pedal point P. Permuting the roles played by the angles  $\alpha, \beta, \gamma$  in (ii), yields a total of six such area-minimizing triangles, which are pedal relative to six pedal points, say,  $P_1, \dots, P_6$ . The main result of the paper is the fact that there exists a circle which contains all six points.

### 1 Introduction

Consider the following question in planar geometry: given three, nonconcurrent lines,  $L_1, L_2, L_3$ , along with three angles  $\alpha$ ,  $\beta$ ,  $\gamma \in (0, \pi)$  satisfying  $\alpha + \beta + \gamma = \pi$ , find three points  $X_1, X_2, X_3$  such that

$$X_1 \in L_1, \ X_2 \in L_2, \ X_3 \in L_3,$$
 (1.1)

and

$$\angle X_2 X_1 X_3 = \alpha, \quad \angle X_1 X_2 X_3 = \beta, \quad \angle X_1 X_3 X_2 = \gamma.$$
 (1.2)

Somewhat informally, with A, B, C playing the roles of the points of mutual intersection of the lines  $L_1, L_2, L_3$ , this asks whether given a fixed triangle ABC, one can inscribe a triangle  $X_1X_2X_3$  of an arbitrarily prescribed shape. For a triangle, "shape" simply signifies the measures of its angles (in the current scenario  $\alpha, \beta, \gamma$ ).

Although not necessarily obvious, this question turns out to have an affirmative answer. In fact, the collection  $\mathcal{I}_{\alpha\beta\gamma}$  of all triangles  $X_1X_2X_3$  satisfying (1.1) and (1.2) has infinite cardinality. Since all triangles in  $\mathcal{I}_{\alpha\beta\gamma}$  have the same shape, what distinguishes them is their size. It is not too difficult to show that  $\mathcal{I}_{\alpha\beta\gamma}$  contains triangles of arbitrarily large areas, hence, the most distinguished triangle in  $\mathcal{I}_{\alpha\beta\gamma}$  is the one of the smallest area, called, for the purposes of this introduction,  $\Delta_{\alpha\beta\gamma}$ . Part of our paper is devoted to studying this minimizer, which enjoys many interesting properties. Among other things we show that this triangle of minimal area is unique, and is also a pedal triangle. Recall that a pedal triangle is a triangle whose vertices are the projections of a point, called the pedal point of the pedal triangle, onto the sides of another triangle. Let  $P_{\alpha\beta\gamma}$  be the pedal point of  $\Delta_{\alpha\beta\gamma}$ .

Of course, similar considerations apply to the situation when the ordered triplet  $(\alpha, \beta, \gamma)$  in (1.2) is replaced by any of its permutations

$$(\alpha, \beta, \gamma), (\alpha, \gamma, \beta), (\beta, \alpha, \gamma), (\beta, \gamma, \alpha), (\gamma, \alpha, \beta), (\gamma, \beta, \alpha).$$
 (1.3)

<sup>\*2000</sup> Math Subject Classification. Primary: 51M25, 51M16. Secondary 51M04, 51M15. Key words: inscribed and circumscribed triangles, pedal triangles, geometrical transformations

As before, these permutations lead to considering the collections  $\mathcal{I}_{\alpha\beta\gamma}$ ,  $\mathcal{I}_{\alpha\gamma\beta}$ , ..., defined as above, which in turn have their own minimizers,

$$\Delta_{\alpha\beta\gamma}, \ \Delta_{\alpha\gamma\beta}, \ \Delta_{\beta\alpha\gamma}, \ \Delta_{\beta\gamma\alpha}, \ \Delta_{\gamma\alpha\beta}, \ \Delta_{\gamma\beta\alpha}.$$
 (1.4)

Once again, they are all pedal triangles with respect to six points, say,

$$P_{\alpha\beta\gamma}, P_{\alpha\gamma\beta}, P_{\beta\alpha\gamma}, P_{\beta\gamma\alpha}, P_{\gamma\alpha\beta}, P_{\gamma\beta\alpha}.$$
 (1.5)

Remarkably, even though these six points are obtained via constructions that start from starkly different sets of premises, they turn out to be concyclic. That is,

**Theorem 1.1** (THE SIX POINT CIRCLE THEOREM). There exists a circle that contains all six points in (1.5).

This is the main result of the paper (cf. Theorem 3.2 for a formal statement). The starting point in its proof is to consider, besides the area-minimizing triangle inscribed in  $\Delta ABC$  and having prescribed angles, the area-maximizing triangle circumscribed to  $\Delta ABC$ , having the same angles. This dual aspect of the problem under consideration is very useful in understanding the nature of the extremal triangles of a prescribed shape. A concrete step in this regard is proving a structure theorem, detailing how these three triangles are related to one another. For instance, we show that the area of the original triangle is the geometric mean of the area-minimizer and the area-maximizer triangles; cf. Theorem 2.4. Since the conclusion in Theorem 1.1 involves a circle, it is not unnatural that another important tool in its proof is the geometrical inversion.

In §2, the properties of the area-minimizing and area-maximizing triangles having prescribed angles and being, respectively, inscribed and circumscribed in a given triangle are examined. Finally, §3 is devoted to presenting the proof of The Six Point Circle Theorem. The presentation in this paper is largely self-contained. For more basic concepts and definitions related to the geometry of the triangle the interested reader is referred to, e.g. [2], and the references therein.

# 2 Extremal Triangles of a Prescribed Shape

Throughout, we let  $\triangle ABC$  denote the triangle with vertices A, B, C and denote by  $|\triangle ABC|$  the area of  $\triangle ABC$ .

Given a triangle  $A_1A_2A_3$  along with a triangle  $B_1B_2B_3$  inscribed in it, we first recall a result from [3] that gives a procedure for obtaining a triangle,  $C_1C_2C_3$ , that is inscribed in  $\Delta B_1B_2B_3$  and is homotopic to  $\Delta A_1A_2A_3$ . By definition, two triangles are homotopic if their corresponding sides are parallel. The concurrence point of the lines passing through the corresponding vertices of two homotopic triangles is the homotopy center. The justification of the following useful result is left to the interested reader (a proof is given in [3]).

**Proposition 2.1.** Let  $\Delta A_1 A_2 A_3$  be arbitrary and assume that  $B_1 \in A_2 A_3$ ,  $B_3 \in A_2 A_1$ ,  $B_2 \in A_1 A_3$  (see Figure 1 below). Take  $C_1 \in B_2 B_3$ ,  $C_2 \in B_1 B_3$ ,  $C_3 \in B_1 B_2$  such that

$$\frac{A_1B_3}{B_3A_2} = \frac{B_2C_3}{C_3B_1}, \quad \frac{A_2B_1}{B_1A_3} = \frac{B_3C_1}{C_1B_2}, \quad \frac{A_3B_2}{B_2A_1} = \frac{B_1C_2}{C_2B_3}, \tag{2.6}$$

Then  $\Delta A_1 A_2 A_3$  and  $\Delta C_1 C_2 C_3$  are homotopic and, in addition,  $|\Delta B_1 B_2 B_3|$  is the geometric mean of  $|\Delta A_1 A_2 A_3|$  and  $|\Delta C_1 C_2 C_3|$ , i.e.

$$|\Delta B_1 B_2 B_3|^2 = |\Delta A_1 A_2 A_3| \cdot |\Delta C_1 C_2 C_3|. \tag{2.7}$$

Conversely, if  $\Delta A_1 A_2 A_3$  and  $B_1 \in A_2 A_3$ ,  $B_3 \in A_2 A_1$ ,  $B_2 \in A_1 A_3$  are given and  $C_1 \in B_2 B_3$ ,  $C_2 \in B_1 B_3$ ,  $C_3 \in B_1 B_2$  are such that  $\Delta A_1 A_2 A_3$  and  $\Delta C_1 C_2 C_3$  are homotopic, then (2.6) and (2.7) hold.




Figure 1

**Definition 2.2.** (i) Call a triangle XYZ inscribed in a given triangle ABC if each of the lines AB, BC and AC contain precisely one of the vertices X, Y, Z. Call  $\Delta XYZ$  circumscribed to  $\Delta ABC$  if  $\Delta ABC$  is inscribed in  $\Delta XYZ$ .

- (ii) Given two fixed triangles ABC, called of reference, and MNP, called fundamental, let  $\mathcal{I}$  be the set of all triangles which are similar to the fundamental one and are inscribed in the triangle of reference, and let  $\mathcal{C}$  be the set of all triangles similar to the fundamental one and circumscribed to the reference one.
- (iii) Given  $\triangle$  ABC, triangle of reference, and a fundamental triangle with angles  $\alpha, \beta, \gamma$ , denote by  $\mathcal{I}_{\alpha\beta\gamma}$  the set of all triangles XYZ with X,Y,Z belonging, respectively, to the lines BC, AC, AB, and for which  $\angle X$ ,  $\angle Y$ ,  $\angle Z$  coincide with  $\alpha, \beta, \gamma$  (in this order).

Likewise, let  $C_{\alpha\beta\gamma}$  denote the set of all triangles XYZ for which A, B, C belonging, respectively, to the lines YZ, XZ, XY, and for which  $\not AX$ ,  $\not AY$ ,  $\not AZ$  coincide with  $\alpha$ ,  $\beta$ ,  $\gamma$  (in this order).

Clearly,

$$\mathcal{I} = \mathcal{I}_{\alpha\beta\gamma} \bigcup \mathcal{I}_{\alpha\gamma\beta} \bigcup \mathcal{I}_{\beta\alpha\gamma} \bigcup \mathcal{I}_{\beta\gamma\alpha} \bigcup \mathcal{I}_{\gamma\alpha\beta} \bigcup \mathcal{I}_{\gamma\beta\alpha}, 
\mathcal{C} = \mathcal{C}_{\alpha\beta\gamma} \bigcup \mathcal{C}_{\alpha\gamma\beta} \bigcup \mathcal{C}_{\beta\alpha\gamma} \bigcup \mathcal{C}_{\beta\gamma\alpha} \bigcup \mathcal{C}_{\gamma\alpha\beta} \bigcup \mathcal{C}_{\gamma\beta\alpha}.$$
(2.8)

Our next result shows the existence of area-minimizers and area-maximizers in the above classes. The issue of uniqueness for these extremal triangles will be dealt with in Theorem 2.4 below.

**Theorem 2.3.** There exists a triangle  $\Delta M_1 N_1 P_1 \in \mathcal{I}_{\alpha\beta\gamma}$  which has the smallest area amongst all the triangles in the class  $\mathcal{I}_{\alpha\beta\gamma}$ . Also, there exists a triangle  $\Delta M_2 N_2 P_2 \in \mathcal{C}_{\alpha\beta\gamma}$  which has the largest area amongst all the triangles in the class  $\mathcal{C}_{\alpha\beta\gamma}$ . In addition, any two such triangles satisfy

$$|\Delta M_1 N_1 P_1| \cdot |\Delta M_2 N_2 P_2| = |\Delta ABC|^2. \tag{2.9}$$

In fact, similar results are valid for each of the classes appearing in (2.8), including  $\mathcal{I}$  and  $\mathcal{C}$ .

*Proof.* It suffices to only deal with the class  $\mathcal{I}_{\alpha\beta\gamma}$  since a similar reasoning applies to all the classes in (2.8). The first issue is to show that there exists at least one area-minimizer in  $\mathcal{I}_{\alpha\beta\gamma}$ . We briefly sketch the argument for this, which is based on results from Calculus and Analytical Geometry. To get started, fix a unit vector  $\vec{w} = w_1 \vec{i} + w_2 \vec{j}$  and consider the class of all triangles XYZ satisfying

It is then clear that the coordinates of Y and Z depend linearly on those of X, with coefficients varying continuously with  $w_1, w_2$ . Since X moves on a line (i.e., BC), it follows that the geometrical locus of the vertex Y is itself a line. In addition, the characteristics of this line, called L, depend continuously on  $w_1, w_2$ . Now, the position of X which, under the constraints (2.10), yields a triangle XYZ in the class  $\mathcal{I}_{\alpha\beta\gamma}$ , is the one corresponding to the case when  $Y = L \cap AC$ . Linear algebra considerations then show that the coordinates of X in this critical case are continuously dependent on  $w_1, w_2$ . Using the formula which expresses the area of a triangle in terms of the coordinates of its vertices, we then deduce that  $|\Delta XYZ|$  -corresponding the case when  $\Delta XYZ \in \mathcal{I}_{\alpha\beta\gamma^-}$  is a continuous function in  $w_1, w_2$ . Since  $w_1$  and  $w_2$  vary in a compact set, it follows that the assignment  $(w_1, w_2) \mapsto |\Delta XYZ|$  is bounded and it attains its extrema. In particular, there exists an area-minimizing triangle in  $\mathcal{I}_{\alpha\beta\gamma}$ .

To prove that there exists (at least) one area-maximizing triangle in  $C_{\alpha\beta\gamma}$ , we proceed as follows. Starting with an area-minimizing triangle  $\Delta M_1 N_1 P_1$  (whose existence has just been established), construct a new triangle,  $\Delta M_2 N_2 P_2$  by taking the parallels through A, B and C to the sides of  $\Delta M_1 N_1 P_1$ . As a result,  $\Delta M_2 N_2 P_2$  is homotopic to  $\Delta M_1 N_1 P_1$  and circumscribed to  $\Delta ABC$ . Hence,  $\Delta M_2 N_2 P_2 \in C_{\alpha\beta\gamma}$  and, thanks to Proposition 2.1,

$$|\Delta M_2 N_2 P_2| \cdot |\Delta M_1 N_1 P_1| = |\Delta ABC|^2. \tag{2.11}$$

We claim that  $\Delta M_2 N_2 P_2$  is area-maximizing in the class  $\mathcal{C}_{\alpha\beta\gamma}$ . Indeed, if  $\Delta M''N''P'' \in \mathcal{C}_{\alpha\beta\gamma}$  is arbitrary, let  $M' \in BA$ ,  $N' \in BC$ ,  $P' \in AC$  (cf. Figure 2) be such that

$$\frac{AM'}{M'B} = \frac{P''C}{CN''}, \quad \frac{BN'}{N'C} = \frac{M''A}{AP''}, \quad \frac{CP'}{P'A} = \frac{N''B}{BM''}.$$
 (2.12)

Proposition 2.1 gives that  $\Delta M'N'P'$  and  $\Delta M''N''P''$  are homotopic (thus  $\Delta M'N'P' \in \mathcal{I}_{\alpha\beta\gamma}$ ) and

$$|\Delta M''N''P''| \cdot |\Delta M'N'P'| = |\Delta ABC|^2. \tag{2.13}$$

Since by definition  $|\Delta M'N'P'| \ge |\Delta M_1N_1P_1|$ , (2.11), (2.13) imply that  $|\Delta M''N''P''| \le |\Delta M_2N_2P_2|$ . This justifies our claim that  $\Delta M_2N_2P_2$  is area-maximizing in the class  $\mathcal{C}_{\alpha\beta\gamma}$ .

Going further, we note that proving (2.9) uses a similar circle of ideas. Concretely, let  $M' \in BA$ ,  $N' \in BC$ ,  $P' \in AC$  (cf. Figure 2) be such that

$$\frac{AM'}{M'B} = \frac{P_2C}{CN_2}, \quad \frac{BN'}{N'C} = \frac{M_2A}{AP_2}, \quad \frac{CP'}{P'A} = \frac{N_2B}{BM_2}.$$
(2.14)

Proposition 2.1 shows that  $\Delta M'N'P'$  and  $\Delta M_2N_2P_2$  are homotopic and

$$|\Delta M_2 N_2 P_2| \cdot |\Delta M' N' P'| = |\Delta ABC|^2. \tag{2.15}$$

The former property also entails  $\Delta M'N'P' \in \mathcal{I}_{\alpha\beta\gamma}$  and, hence,

$$|\Delta M_1 N_1 P_1| \le |\Delta M' N' P'|. \tag{2.16}$$




Figure 2

The parallels through A, B and C to the sides of the triangle  $\Delta M_1 N_1 P_1$  determine the triangle M''N''P'' which is homotopic to  $\Delta M_1 N_1 P_1$  and circumscribed to  $\Delta ABC$ . As a result,  $\Delta M''N''P'' \in \mathcal{C}_{\alpha\beta\gamma}$  which forces

$$|\Delta M''N''P''| \le |\Delta M_2 N_2 P_2|. \tag{2.17}$$

In this context, Proposition 2.1 also gives that

$$|\Delta M''N''P''| \cdot |\Delta M_1 N_1 P_1| = |\Delta ABC|^2. \tag{2.18}$$

By combining (2.15)-(2.18) we obtain

$$|\Delta ABC|^2 = |\Delta M_1 N_1 P_1| |\Delta M'' N'' P''|$$
  
 $\leq |\Delta M_2 N_2 P_2| |\Delta M' N' P'| = |\Delta ABC|^2,$  (2.19)

so that, we actually have  $|\Delta M''N''P''| = |\Delta M_2N_2P_2|$ ,  $|\Delta M_1N_1P_1| = |\Delta M'N'P'|$ . Therefore,  $|\Delta ABC|^2 = |\Delta M_1N_1P_1||\Delta M''N''P''| = |\Delta M_1N_1P_1||\Delta M_2N_2P_2|$ , finishing the proof of (2.9).

**Remark 1.** It is implicit in the proof of the above theorem that if  $\Delta M_1 N_1 P_1$  is an area-minimizer in the class  $\mathcal{I}_{\alpha\beta\gamma}$ , then the triangle  $M_2 N_2 P_2$  that is in  $\mathcal{C}_{\alpha\beta\gamma}$  and is homotopic to  $\Delta M_1 N_1 P_1$  (constructed as in Proposition 2.1) is an area-maximizer in the class  $\mathcal{C}_{\alpha\beta\gamma}$ . Conversely, if  $\Delta M_2 N_2 P_2$  is an area-maximizer in the class  $\mathcal{C}_{\alpha\beta\gamma}$ , then the triangle  $M_1 N_1 P_1$  that is in  $\mathcal{I}_{\alpha\beta\gamma}$  and is homotopic to  $\Delta M_2 N_2 P_2$  (again, constructed as in Proposition 2.1) is an area-minimizer in the class  $\mathcal{I}_{\alpha\beta\gamma}$ .

The main result in this section describes the structure of the configuration consisting of a given triangle ABC along with the area-minimizing triangle in  $\mathcal{I}_{\alpha\beta\gamma}$  and the area-maximizing triangle in  $\mathcal{C}_{\alpha\beta\gamma}$ . Recall that an antipedal triangle of a point P with respect to  $\Delta ABC$  is the triangle whose vertices are the intersection points of the perpendiculars to PA, PB, and PC, passing through A, B, and C, respectively. Also, the isogonal of a point P in  $\Delta ABC$  is the concurrence point of the reflections of the lines PA, PB, and PC across the angle bisectors of A, B, and C.

**Theorem 2.4.** Retain notation and conventions introduced earlier. Given  $\triangle ABC$ , reference triangle, and  $\triangle MNP$  fundamental triangle with angles  $\alpha, \beta, \gamma \in (0, \pi)$ , let  $\mathcal{I}_{\alpha\beta\gamma}$  and  $\mathcal{C}_{\alpha\beta\gamma}$  be as before.

Then there exist a unique  $\Delta EFD \in \mathcal{I}_{\alpha\beta\gamma}$  which is area-minimizing in this class, along with a unique  $\Delta QRS \in \mathcal{C}_{\alpha\beta\gamma}$  which is area-maximizing in this class. In addition, these triangles enjoy the following properties:

- (i) The triangles EFD and QRS are homotopic;
- (ii) The triangles EFD and QRS, are the pedal and the antipedal triangles of some points K and L, respectively, with respect to  $\Delta ABC$ ;
- (iii) Let T be the concurrence point of the projections of Q, R, S on the sides of the triangle ABC. Then the pairs of points (L,K) and (T,L) are isogonal in the triangles ABC and QRS, respectively;
- (iv) If O is the homotopy center of  $\Delta EFD$  and  $\Delta QRS$ , then O, K, T lie on the same line.

Before proceeding with the proof of this theorem, a comment is in order.

**Remark 2.** As a corollary of the above theorem, there exists a unique area-minimizing triangle in each of the classes  $\mathcal{I}$ ,  $\mathcal{I}_{\alpha\beta\gamma}$ , ..., listed in the first line of (2.8). Moreover, there exists a unique area-maximizing triangle in each of the classes  $\mathcal{C}$ ,  $\mathcal{C}_{\alpha\beta\gamma}$ , ..., listed in the second line of (2.8). In each case, similar properties to (i) – (iv) above hold.




Figure 3

Since  $\not AM + \not AN + \not AP = \not AQ + \not AR + \not AS = \pi$ , it is not difficult to see that these three circles, have a common point, which we denote by L. Observe next that a particular position of the point S on the circle  $(O_1)$  determines also the positions of the points Q and R on the circles  $(O_2)$  and  $(O_3)$ , respectively. The idea is now to determine the position of the point S on the circle  $(O_1)$  for which the triangle QRS has maximal area.




Figure 4

Since the angles of  $\Delta QRS$  are fixed, maximizing  $|\Delta QRS|$  amounts to maximizing the length of one of its sides, say SQ. Let  $O_1'$  and  $O_2'$  be, respectively, the projections of  $O_1$  and  $O_2$  onto the line SQ, and let  $O_1'' \in O_1'O_1$  be such that  $O_1''O_2' \parallel O_1O_2$ . Then we can write

$$O_1O_2 = O_1''O_2' \ge O_1'O_2' = BO_1' + BO_2' = \frac{BS}{2} + \frac{BQ}{2} = \frac{SQ}{2}.$$
 (2.20)

This implies  $2O_1O_2 \geq SQ$ , an inequality from which we see that the maximum of  $|\Delta QRS|$  is attained when  $SQ \parallel O_1O_2$ .

Since the location of the points  $O_1$ ,  $O_2$ ,  $O_3$  depends only on  $\triangle ABC$  and angles  $\alpha, \beta, \gamma$ , the result just established shows that there exists a unique area-maximizing triangle in  $\mathcal{C}_{\alpha\beta\gamma}$ . Given an area-minimizing triangle  $\triangle DEF$  in the class  $\mathcal{I}_{\alpha\beta\gamma}$ , we can associate with it a homotopic triangle  $\triangle QRS$  in  $\mathcal{C}_{\alpha\beta\gamma}$ , which can be constructed following the recipe given in Proposition 2.1. Using the Remark 1, the latter triangle is area-maximizing in  $\mathcal{C}_{\alpha\beta\gamma}$  and, hence, must coincide with the unique area-maximizing triangle described in the first part of the current proof. This shows that  $\triangle DEF$  is itself uniquely determined by the property that it is area-minimizing in  $\mathcal{I}_{\alpha\beta\gamma}$ . This concludes the proof of the first claim made in the statement of Theorem 2.4. The above considerations also show that  $\triangle DEF$  and  $\triangle QRS$  are homotopic, thus proving item (i).

As regards (ii), let us recall from the first part of the proof that the triangle QRS has a maximal area when  $S = LO_1 \cap (O_1)$ ,  $Q = LO_2 \cap (O_2)$ ,  $R = LO_3 \cap (O_3)$ . In particular, this implies that the triangles QRS and  $O_1O_2O_3$  are homotopic with homotopy center L and dilation factor  $\frac{1}{2}$ . Next, since  $O_1O_2 \parallel SQ$  and  $O_1O_2 \perp BL$ , it follows that  $BL \perp SQ$ . Similarly  $LC \perp RQ$  and  $LA \perp SR$ . Altogether, these observations imply that the triangle SRQ is the antipedal of L with respect to the triangle ABC. Now, using that  $DF \parallel SR$ ,  $DE \parallel SQ$ , and  $FE \parallel RQ$  we obtain that  $AL \perp DF$ ,  $BL \perp DE$ , and  $CL \perp EF$ . This means that the projections of the vertices of  $\Delta ABC$  on the sides of  $\Delta DEF$  are concurrent, therefore establishing that the triangles ABC and DEF are orthologic. As is well-know, this gives that the perpendiculars from D, E, F onto AB, BC, AC respectively, are also concurrent in a point, say K. Hence,  $\Delta DEF$  is the pedal triangle of K with respect to the triangle ABC. This finishes the proof of (ii).

Consider next (iii). Since the triangles ABC and SQR are orthologic, there exists a point T such that  $ST \perp AB$ ,  $QT \perp BC$ , and  $RT \perp AC$ . In the quadrilateral ADKF we have

Therefore,

$$\angle FDK = \angle LAD.$$
 (2.22)

From (2.21) and (2.22) we see that the lines AL and AK are isogonal. Similarly, BL and BK are isogonal, proving that L and K are isogonal points in  $\Delta ABC$ . This concludes the proof of (iii).

To prove (iv), we start by observing that the dilation of center O and factor  $\frac{SQ}{DE}$  (i.e., the dilation which takes  $\Delta DEF$  into  $\Delta SQR$ ) transforms the lines DK, EK and FK into the lines ST, RT and QT, respectively. (Indeed, this is a direct corollary of the fact that dilations map parallel lines into parallel lines.) Thus, under such a transformation, the point K is mapped into T and, as a consequence, the points O, K and T are collinear. The proof of Theorem 2.4 is now finished.  $\Box$ 

The result below essentially says the that a pedal triangle (relative to  $\Delta ABC$ ) is area-minimizing amongst all triangles of a similar shape inscribed in  $\Delta ABC$ .

**Theorem 2.5.** Let P be a point contained inside the circumcircle of a given triangle ABC. Let D, E, F be the projections of P onto the sides BC, AC and AB, respectively, and set

$$\alpha := \angle EDF, \quad \beta := \angle DEF, \quad \gamma := \angle EFD. \tag{2.23}$$

Then  $\Delta DEF$  is the area-minimizing triangle in the class  $\mathcal{I}_{\alpha\beta\gamma}$  (cf. (ii) of Definition 2.2).

Proof. Let K be the isogonal of P (relative to  $\triangle ABC$ ) and denote by  $\triangle QRS$  the antipedal of K with respect to  $\triangle ABC$ . Since  $\triangle DEF$  is the pedal triangle of the point P with respect to  $\triangle ABC$ , then quadrilateral AFPE is inscribable. This implies that  $\angle PEF + \angle PAF = \frac{\pi}{2}$ . Next, since K is the isogonal of P, this further entails  $\angle PEF + \angle KAE = \frac{\pi}{2}$ . This means that  $AK \perp EF$ , i.e.  $RS \parallel EF$ . Via a similar reasoning, we can show that the two other pairs of corresponding sides in  $\triangle QRS$  and  $\triangle DEF$  are parallel. This shows that  $\triangle QRS$  and  $\triangle DEF$  are homotopic. In particular, the angles of  $\triangle QRS$  are  $\alpha$ ,  $\beta$ ,  $\gamma$ .

Recall the (three, concurrent) circles  $(O_1)$ ,  $(O_2)$ ,  $(O_3)$  introduced in the proof of Theorem 2.4. Then  $S \in (O_1)$ ,  $R \in (O_2)$ ,  $Q \in (O_3)$ . Given that the quadrilateral AKCR is inscribable, it follows that  $K \in (O_1)$ . Similarly,  $K \in (O_2)$ ,  $K \in (O_3)$ . Hence, K is the common point of  $(O_1)$ ,  $(O_2)$ ,  $(O_3)$ , which implies that  $\Delta QRS$  is homotopic to  $\Delta O_1O_2O_3$ . As pointed out already in the proof of Theorem 2.4, this necessarily implies that  $\Delta QRS$  is a maximizer in the class  $\mathcal{C}_{\alpha\beta\gamma}$ . In turn, by (iii) in Theorem 2.4 this forces  $\Delta DEF$  to be an area minimizer in  $\mathcal{I}_{\alpha\beta\gamma}$ .

We end with a result which appears to be folklore (see [1]; a new proof has been given in [3]).

**Theorem 2.6.** Assume ABC to be a given triangle and denote by O and R the center and the radius of the circumcircle, respectively. Let M be an arbitrary point in  $\triangle ABC$  and let  $\triangle \alpha \beta \gamma$  be the pedal triangle of M with respect to  $\triangle ABC$ . Then

$$\frac{|\Delta\alpha\beta\gamma|}{|\Delta ABC|} = \frac{|R^2 - OM^2|}{4R^2}.$$
 (2.24)

#### 3 The Main Result

This section is devoted to stating and proving the main result in this paper, Theorem 3.2. First, we take care of a number of prerequisites, starting with the result below.

**Theorem 3.1.** Let M be a point contained in the circumcircle of  $\triangle ABC$ , which is assumed to have center O and radius R. In addition, let  $N \in OM$  be such that  $OM \cdot ON = R^2$ . Then the pedal triangles,  $\triangle M_1 M_2 M_3$  and  $\triangle N_1 N_2 N_3$ , of the points M and N with respect to  $\triangle ABC$  are similar (with pairs of equal angles having vertices located on the same sides of  $\triangle ABC$ ). Furthermore,

$$|\Delta M_1 M_2 M_3| \le |\Delta N_1 N_2 N_3|. \tag{3.25}$$

Conversely, if M and  $N \in OM$  are two points such that their pedal triangles with respect to  $\Delta ABC$  are similar (with pairs of equal angles having vertices on the same sides of  $\Delta ABC$ ), then  $OM \cdot ON = R^2$ .

*Proof.* Let X, Y and Z be the centers of the circles AMB, BMC and AMC, respectively (see Figure 5). By an inversion of center O and modulus R, the triangle ABC and the circle (O) remain unchanged. The circles (X), (Y) and (Z) are transformed in the circles (X'), (Y') and (Z'), respectively, passing through A and B, B and C, A and C, respectively, and have a common point N, which is the inverse transform of M (see Figure 6).




Figure 5




Figure 6




Figure 7

$$\angle BNA = \angle BMA - 2\angle ACB.$$
 (3.26)

If  $M_1M_2M_3$  and  $N_1N_2N_3$  are the pedal triangles of M and N respectively, with

$$M_1, N_1 \in AB, M_2, N_2 \in BC, M_3, N_3 \in AC,$$
 (3.27)

then

In addition, the quadrilaterals  $N_3NN_1A$  and  $N_2NN_1B$  can be inscribed in a circle, therefore

$$\angle N_1 N A = \angle N_1 N_3 A \text{ and } \angle N_1 N_2 B = \angle B N N_1.$$
 (3.29)

Furthermore,

$$\angle BNA = \angle N_1NA + \angle BNN_1, \text{ and } \angle N_1 = \pi - \angle N_2 - \angle N_3,$$
 (3.30)

Combining (3.26), (3.28), (3.29), and (3.30), we arrive at the conclusion that

$$\angle N_1 = \angle BNA + \angle ACB.$$
 (3.31)

It is also possible to write

$$\angle BMA = \pi - \angle MAM_1 - \angle MBM_1, \tag{3.32}$$

$$\angle ACB = \pi - \angle CAB - \angle CBA,$$
 (3.33)

$$\angle CAB = \angle MAC + \angle MAM_1, \tag{3.34}$$

$$\angle CBA = \angle MBC + \angle MBM_1. \tag{3.35}$$

Therefore, by combining (3.32)-(3.35), we obtain

$$\angle BMA - \angle ACB = \angle MAC + \angle MBC.$$
 (3.36)

Since the quadrilaterals  $M_3AM_1M$  and  $MM_1BM_2$  can be inscribed in a circle,

$$\angle MAC = \angle MM_1M_3 \text{ and } \angle MBC = \angle MM_1M_2.$$
 (3.37)

However,

so therefore, by (3.36), (3.37), and (3.38),

$$\angle M_1 = \angle BMA - \angle ACB.$$
 (3.39)

The identities (3.39) and (3.31) imply that  $\not \perp N_1 = \not \perp M_1$ . A similar reasoning can be used to show that  $\not \perp N_2 = \not \perp M_2$ , and that  $\not \perp N_3 = \not \perp M_3$ , proving that  $\Delta N_1 N_2 N_3$  and  $\Delta M_1 M_2 M_3$  are similar, with their corresponding angles' vertices located on the same sides of the triangle ABC.

Next we will prove (3.25). By Theorem 2.6,

$$\frac{|\Delta M_1 M_2 M_3|}{|\Delta N_1 N_2 N_3|} = \frac{R^2 - OM^2}{ON^2 - R^2}$$
(3.40)

and the last fraction is  $\leq 1$ , as it can be seen from the fact that  $OM \leq R \leq ON$  and  $OM \cdot ON = R^2$ . Hence, (3.25) follows.

Now we will prove the converse statement referred to in Theorem 3.1. In order for  $\Delta N_1 N_2 N_3$  and  $\Delta M_1 M_2 M_3$  to be similar, it is a necessary condition that  $\not \Delta BNA = \not \Delta BMA - 2\not \Delta ACB$ , which is equivalent to  $\not \Delta BNA = \frac{\widehat{f}}{2} - \frac{\widehat{g}}{2}$ . This can only occur when  $\not \Delta BNA = \frac{\widehat{h}}{2}$  which implies that N is on the circle X'. A similar reasoning can be applied to show that N must be on the circles Y' and Z', proving that N is the inversion of M. This shows that the converse statement is also valid, completing the proof of Theorem 3.1.

- Remark 3. (i) From the reasoning above we see that there exist precisely six points in the interior of the circumscribed circle of  $\Delta ABC$  such that their antipedal triangles with respect to  $\Delta ABC$  are similar to a given reference triangle MNP. We shall call these points the interior points of  $\Delta MNP$  with respect to  $\Delta ABC$ .
- (ii) Likewise, there exist precisely six points in the exterior of the circumcircle of  $\Delta ABC$  with the same property as above. We shall call these points the exterior points of  $\Delta MNP$  with respect to  $\Delta ABC$ .

After this preamble, we are ready to state and prove our main result:

**Theorem 3.2.** The pedal points of the six triangles,  $\Delta_{\alpha\beta\gamma}$ ,  $\Delta_{\alpha\gamma\beta}$ ,  $\Delta_{\beta\alpha\gamma}$ ,  $\Delta_{\beta\gamma\alpha}$ ,  $\Delta_{\gamma\alpha\beta}$ ,  $\Delta_{\gamma\beta\alpha}$ , that are area-minimizers in the classes  $\mathcal{I}_{\alpha\beta\gamma}$ ,  $\mathcal{I}_{\alpha\gamma\beta}$ ,  $\mathcal{I}_{\beta\alpha\gamma}$ ,  $\mathcal{I}_{\beta\gamma\alpha}$ ,  $\mathcal{I}_{\gamma\alpha\beta}$ ,  $\mathcal{I}_{\gamma\beta\alpha}$ , respectively, all lie on a circle.

Proving Theorem 3.2 is equivalent to proving the following. Consider ABC, triangle of reference, and  $\Delta D_0 E_0 F_0$  a fixed fundamental triangle (in the terminology of Definition 2.2).

Let  $M_i$ , i = 1, 2, ..., 6, be six points such that, for each  $i \in \{1, 2, ..., 6, \}$ , the pedal triangle of  $M_i$  with respect to  $\Delta ABC$  is  $\Delta D_i E_i F_i$  with

and

$$D_1, D_2, E_3, E_4, F_5, F_6 \in AB,$$

$$D_3, D_5, E_1, E_6, F_2, F_4 \in BC,$$

$$D_4, D_6, E_2, E_5, F_1, F_3 \in AC,$$

$$(3.42)$$

(see Figure 8). Then the points  $M_i$ , i = 1, 2, ..., 6, lie on the same circle.




Figure 8

*Proof.* Since  $\Delta D_1 E_1 F_1$  is the antipedal triangle of  $M_1$ , with respect to  $\Delta ABC$ , we can obtain (reasoning similarly to what was done in the proof of Theorem 3.1 to show that  $\not AM_1 = \not AMA - \not ACB$ ) that

Similarly we obtain  $\not \Delta D_2 = \not \Delta B M_2 A - \not \Delta C$ . From  $\not \Delta D_0 = \not \Delta D_1 = \not \Delta D_2$ , it follows that  $\not \Delta B M_1 A = \not \Delta B M_2 A$ , i.e.,

$$M_1, M_2$$
 lie on a circle  $(O_D)$ , passing through  $A$  and  $B$  and with the property that  $\frac{1}{2}\overrightarrow{AB} = \cancel{4}AM_1B = \cancel{4}C + \cancel{4}D_0$ . (3.44)

Likewise (see Figure 9),

$$M_3, M_4$$
 lie on a circle  $(O_E)$ , passing through  $A$  and  $B$  and with the property that  $\frac{1}{2}\overrightarrow{AB} = \cancel{\bot}C + \cancel{\bot}E_0$ , (3.45)

$$M_5, M_6$$
 lie on a circle  $(O_F)$ , passing through  $A$  and  $B$  and with the property that  $\frac{1}{2}\overrightarrow{AB} = \cancel{\bot}C + \cancel{\bot}F_0$ , (3.46)

$$M_6, M_4$$
 lie on a circle  $(O'_D)$ , passing through  $A$  and  $C$  and with the property that  $\frac{1}{2}\overrightarrow{AC} = \cancel{4}B + \cancel{4}D_0$ , (3.47)

$$M_5, M_2$$
 lie on a circle  $(O_E')$ , passing through  $A$  and  $C$  and with the property that  $\frac{1}{2}\widehat{AC} = \cancel{4}B + \cancel{4}E_0$ , (3.48)

$$M_1, M_3$$
 lie on a circle  $(O'_F)$ , passing through  $A$  and  $C$  and with the property that  $\frac{1}{2}\widehat{AC} = \cancel{\bot}B + \cancel{\bot}F_0$ . (3.49)

The key step in the proof is to make at this stage an inversion of center A and arbitrary modulus. Since the lines AB and AC pass through A, they will remain unchanged. The points B, C become B' and C', respectively. The circles  $(O_D)$ ,  $(O_E)$ ,  $(O_F)$  will be transformed into the lines  $d_D$ ,  $d_E$  and  $d_F$  respectively, which are concurrent at B'. Meanwhile, the circles  $(O'_D)$ ,  $(O'_E)$  and  $(O'_F)$  become the lines  $d'_D$ ,  $d'_E$  and  $d'_F$ , respectively, which are concurrent at C'. The points  $M_1$ ,  $M_2$ ,  $M_3$ ,  $M_4$ ,  $M_5$ ,  $M_6$  will be transformed into  $M'_1$ ,  $M'_2$ ,  $M'_3$ ,  $M'_4$ ,  $M'_5$  and  $M'_6$ , respectively, and  $M'_1$ ,  $M'_2 \in d_D$ ,  $M'_3$ ,  $M'_4 \in d_E$ ,  $M'_5$ ,  $M'_6 \in d_F$ ,  $M'_6$ ,  $M'_4 \in d'_D$ ,  $M'_5$ ,  $M'_2 \in d'_E$ ,  $M'_1$ ,  $M'_3 \in d'_F$ .




Figure 9

The goal is to show that the points  $M_1'$ ,  $M_2'$ ,  $M_3'$ ,  $M_4'$ ,  $M_5'$ ,  $M_6'$  lie on the same circle. We note that since under an inversion the angle between two curves is preserved, we have  $\not \perp (d_D, d_F') = \not \perp ((O_D), (O_F'))$ , as the angle between two circles is defined to be the angle between the tangents at one of their common points. Therefore,  $\not \perp (d_D, d_F') = \not \perp (\alpha A\beta) = \pi - \not \perp (O_DAO_F')$ . However,

$$\cancel{\Delta}(O_D A O_F') = \cancel{\Delta}(O_D A B) + \cancel{\Delta} A + \cancel{\Delta}(C A O_F').$$
(3.50)

On the other hand  $\angle(O_D AB) = \frac{\pi - \cancel{\downarrow}(AO_D B)}{2} = \cancel{\downarrow}C + \cancel{\downarrow}D_0 - \frac{\pi}{2} \text{ and } \cancel{\downarrow}(CAO_F') = \cancel{\downarrow}F_0 + \cancel{\downarrow}B - \frac{\pi}{2}.$ Hence,

$$\measuredangle(O_D A O_F') = (\measuredangle A + \measuredangle B + \measuredangle C) + \measuredangle D_0 + \measuredangle F_0 - \pi = \measuredangle D_0 + \measuredangle F_0, \tag{3.51}$$

and  $\not \Delta(d_D, d_F') = \pi - \not \Delta(O_D A O_F') = \not \Delta E_0$ .

Let  $B'\delta$  be the tangent at B to the circle determined by the points A', B', C'. Then  $\not \perp (d_E, B'C') = \not \perp (\delta B'C') - \not \perp (\delta B', d_E)$ . However,  $\not \perp (\delta B'C') = \not \perp A$ , and  $\not \perp (\delta B', d_E) = \not \perp (C(A, B', C'), d_E) = \not \perp (BC, (O_E)) = -\not \perp B + (\pi - \not \perp C - \not \perp E_0)$ , where C(A, B, C) denotes the circle containing A, B, C. Therefore

We have the pairs of lines  $(d_D, d'_D)$ ,  $(d_E, d'_E)$ ,  $(d_F, d'_F)$  having the same slope as the line BC, and by symmetry, we see that  $M'_4M'_2 \parallel M'_6M'_1 \parallel M'_5M'_3 \parallel B'C'$ . Hence,  $\not \perp (M'_4M'_2B') = \not \perp (M'_2B'C')$  and  $\not \perp (B'M'_3C') = \pi - \not \perp (M'_3B'C') - \not \perp (M'_3C'B') = \pi - \not \perp E_0 - \not \perp F_0 = \not \perp D_0$ . As a consequence, the quadrilateral  $M'_4M'_2M'_1M'_3$  is inscribable. Since  $\not \perp (B'M'_5C') = \not \perp (B'M'_3C') = \not \perp D_0$ , it follows that the quadrilateral  $M'_4M'_2M'_1M'_6$  is also inscribable. The quadrilateral  $M'_4M'_2M'_1M'_6$  is an isosceles trapezoid, hence it is inscribable as well. As a result,  $M'_6$  lies on the circumscribed circle to the quadrilateral  $M'_3M'_1M'_2M'_4$ . Since  $M'_5$  lies on the circumscribed circle to the quadrilateral  $M'_6M'_1M'_3M'_5$ , it follows that all the points  $M_i$ , i = 1, 2, ..., 6, lie on the same circle.

**Remark 4.** Since the interior points lie on a circle, it follows that the exterior points lie on a circle (the transformed under the inversion of center O and modulus R of the former circle).

#### References

- [1] W. Gallatly, The Modern Geometry of the Triangle, Francis Hodgson, London, 1910.
- [2] C. Kimberling, Triangle Centers and Central Triangles, Congressus Numerantium 129, 1998, pp. 295.
- [3] A. Mitrea, On the area of pedal triangles, preprint, (2008).

California Institute of Technology MSC 700, Pasadena, CA 91126, USA

e-mail: amitrea@caltech.edu